

Purdue University
Purdue e-Pubs

LARS Symposia

Laboratory for Applications of Remote Sensing

1-1-1976

Comparison of Vegetation Classes in the Great Dismal Swamp Using Two Individual Landsat Images and a Temporal Composite

Patricia T. Gammon

Virginia P. Carter

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Gammon, Patricia T. and Carter, Virginia P., "Comparison of Vegetation Classes in the Great Dismal Swamp Using Two Individual Landsat Images and a Temporal Composite" (1976). *LARS Symposia*. Paper 139.
http://docs.lib.purdue.edu/lars_symp/139

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

June 29 - July 1, 1976

The Laboratory for Applications of
Remote Sensing

Purdue University
West Lafayette
Indiana

IEEE Catalog No.
76CH1103-1 MPRSD

Copyright © 1976 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

COMPARISON OF VEGETATION CLASSES IN THE
GREAT DISMAL SWAMP USING TWO INDIVIDUAL
LANDSAT IMAGES AND A TEMPORAL COMPOSITE

Patricia T. Gammon
Dismal Swamp National Wildlife Refuge
Suffolk, Virginia

Virginia P. Carter
U. S. Geological Survey
Reston, Virginia

ABSTRACT

The Great Dismal Swamp on the Virginia/North Carolina border has highly diverse vegetation, largely due to past alteration by lumbering, fire and drainage. Vegetation maps are presently being completed using National Aeronautics and Space Administration high and low altitude color infrared photographs and 1:100,000 and 1:24,000 scale black and white orthophoto bases. Having mapped the vegetation to the required precision for management needs, we are now evaluating the application potential of Landsat data to provide a cost-effective, repetitive information source for future monitoring and map revision.

Three data sets were prepared for analysis using two Landsat scenes (Image #1583-15100, February, 1974; Image #1637-15085, April, 1974) which were geometrically corrected, resampled and temporally registered by the Jet Propulsion Laboratory. All four bands were used to analyze individual dates; bands 5 and 7 for each date were used to analyze the temporal composite.

Each of the three data sets provides information on specific aspects of vegetation classification. In addition, comparison of the individual dates and the temporal composite allows further interpretation beyond the information provided by individual analyses.

Using February data, three evergreen classes may be separated; pine, Atlantic white cedar and the dense evergreen shrub community. Deciduous classes may be divided only on the basis of the ground cover: heavy evergreen understory, sparse (mostly deciduous) understory or standing water. No deciduous canopy species can be identified with February data.

Using April data, the boundaries of the pine and cedar classes become less distinct. The evergreen shrub class remains well-defined. Deciduous canopy species may be separated in areas where one species predominates.

By comparing the April and February data, stands of cypress can be identified. The overlap between pine and evergreen understory in February data can be resolved in April by leaf-out of deciduous canopy.

The temporally registered data gives classes much closer in areal extent and location to the classes mapped at 1:100,000 scale from color infrared photographs. The area classified as deciduous over standing water is almost identical to that mapped as cypress/tupelo. The pine, cedar and evergreen shrub categories separate extremely well. A large deciduous class can be identified as mostly maple with some yellow poplar, sweet gum and tupelo. In addition, a class consisting of broad-leaved evergreen shrubs, trees and vines could only be separated using the temporal data.

No single data set provided adequate information for vegetative classification in the Great Dismal Swamp. Seasonal and temporal comparisons provide approximately 70% of the desired precision in vegetative boundary determination and class separation.